

A work done by an isotropic vector force field along an isotropic curve

D. Razpopov and G. Dzhelepov

Abstract. We consider a 4-dimensional Riemannian manifold M with a metric g and an endomorphism Q , whose fourth power is the identity and Q acts as an isometry on g . An associated metric \tilde{g} on (M, g, Q) is determined by both structures g and Q . The metric \tilde{g} is necessary indefinite and it induces isotropic vectors in the tangent space at an arbitrary point on M . The physical forces are represented by vector fields. We study forces whose vectors are in a single tangent space on (M, g, Q) . We calculate the corresponding physical work done by arbitrary forces along arbitrary curves with respect to \tilde{g} . Mainly, we suppose that the vector force fields are isotropic and they act along isotropic curves. We calculate the physical work done by such forces.

M.S.C. 2010: 53C15, 53B20, 53Z05.

Key words: Riemannian manifold; indefinite metric tensor; isotropic vectors; null curves.

1 Introduction

The theory of differentiable manifolds with additional structures have many applications in mathematics and physics. The physical forces on curves of differentiable manifolds are associated with vector fields on the manifolds and then one could find the physical work done by such forces. If a vector force field acts along a curve from one point to another point, the work done by the force is the product of the force and the displacement. For example, with vector fields are modeled forces such as the magnetic and gravitational forces. We consider vector force fields along curves of a Riemannian manifold equipped with an additional indefinite metric. In particular, the forces and curves are described by isotropic (null, light-like) vectors. The properties of null curves of semi-Riemannian manifolds and their important applications in general relativity are studied in detail ([8, 9]). There are some papers concerning physical results on light-like objects of differentiable manifolds (see [1, 2, 11, 12, 13, 15]).

In the present paper we consider a 4-dimensional Riemannian manifold M with a metric g and a tensor Q of type $(1, 1)$, whose fourth power is the identity. Moreover,

Q acts as an isometry on g . Such a manifold (M, g, Q) is defined in [16] and it is studied also in [3, 4, 5, 7, 10]. We consider an associated metric \tilde{g} , introduced in [4], which is necessary indefinite. Therefore, \tilde{g} determines isotropic (null) vectors in every tangent space T_pM at a point p on (M, g, Q) . We investigate the physical work done by arbitrary vector force fields along arbitrary curves in T_pM and in subspaces of T_pM , with respect to \tilde{g} . Especially, we obtain the physical work done by isotropic vector force fields along isotropic curves.

2 Preliminaries

Let M be a 4-dimensional Riemannian manifold with a metric g and a tensor Q of type $(1, 1)$. Let the local coordinates of Q with respect to some coordinate system form the following circulant matrix:

$$(2.1) \quad (Q_i^j) = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \end{pmatrix}.$$

Then Q has the property

$$(2.2) \quad Q^4 = \text{id}.$$

We assume that g is a positive definite metric on M , which satisfies the equality

$$(2.3) \quad g(Ql, Qv) = g(l, v).$$

Remark 2.1. The notations l, v in (2.3) and furthermore r, e_1, e_2, e_3 will stand for arbitrary vector fields or arbitrary vectors in T_pM .

The manifold (M, g, Q) , determined by (2.1) and (2.3), is introduced in [16].

Bearing in mind (2.3), for the angles $\varphi = \angle(v, Qv)$ and $\phi = \angle(v, Q^2v)$ we have

$$(2.4) \quad \cos \varphi = \frac{g(v, Qv)}{g(v, v)} \quad \cos \phi = \frac{g(v, Q^2v)}{g(v, v)}$$

In [16], for (M, g, Q) and for an arbitrary nonzero vector v in T_pM , it is verified that if v induces a Q -basis of T_pM , that is a basis of the type $\{v, Qv, Q^2v, Q^3v\}$, then the angles ϕ and φ satisfy inequalities

$$(2.5) \quad 0 < \phi < \pi, \quad \frac{\phi}{2} < \varphi < \pi - \frac{\phi}{2}.$$

The associated metric \tilde{g} on (M, g, Q) , determined by

$$(2.6) \quad \tilde{g}(l, v) = g(l, Q^2v)$$

is necessary indefinite [4].

A nonzero vector v in T_pM is isotropic with respect to \tilde{g} if

$$(2.7) \quad \tilde{g}(v, v) = 0.$$

An isotropic curve $c : r = r(t)$, where $t \in [\alpha, \beta] \subset \mathbb{R}$, is this one whose tangent vector field dr is isotropic, i.e.

$$(2.8) \quad \tilde{g}(dr, dr) = 0.$$

If F is a vector force field, then the work A done by a force F , with respect to \tilde{g} , moving along a curve c is given by

$$(2.9) \quad A = \int_c \tilde{g}(F, dr),$$

where dr is the elementary displacement on the curve c .

Theorem 2.1. *Let F be an isotropic vector force field and let c be an isotropic curve. If F and the tangent vector field dr of c are linearly dependent, then the work A done by a force F , with respect to \tilde{g} , along the curve c is zero.*

Proof. We have that $F = kdr$, where k is a function. Then, using (2.8) and (2.9), we get $dA = \tilde{g}(kdr, dr) = k\tilde{g}(dr, dr) = 0$, which implies $A = 0$. \square

Due to Theorem 2.1 and having in mind that every vector field and the tangent vector field to his trajectory are linearly dependent we state

Remark 2.2. If c is a trajectory of isotropic force F , then c is also isotropic and the corresponding work A is zero.

3 The work in T_pM

An orthonormal Q -basis $\{v, Qv, Q^2v, Q^3v\}$ exists in every tangent space T_pM on (M, g, Q) ([16]).

We assume that p_{xyzu} is a coordinate system such that the vectors v, Qv, Q^2v and Q^3v lie on the axes p_x, p_y, p_z and p_u , respectively. So p_{xyzu} is an orthonormal coordinate system.

A vector force field F is determined by

$$(3.1) \quad \begin{aligned} F(x, y, z, u) = & P(x, y, z, u)v + R(x, y, z, u)Qv \\ & + S(x, y, z, u)Q^2v + L(x, y, z, u)Q^3v, \end{aligned}$$

where $P = P(x, y, z, u)$, $R = R(x, y, z, u)$, $S = S(x, y, z, u)$, $L = L(x, y, z, u)$ are smooth functions.

A smooth curve c is determined by

$$(3.2) \quad c : r(t) = x(t)v + y(t)Qv + z(t)Q^2v + u(t)Q^3v,$$

where $t \in [\alpha, \beta] \subset \mathbb{R}$.

Theorem 3.1. *Let F be an arbitrary vector force field and let c be an arbitrary smooth curve in T_pM , determined by (3.1) and (3.2). Then the work A done by a force F , with respect to \tilde{g} , along c is*

$$(3.3) \quad A = \int_{\alpha}^{\beta} (Pz'(t) + Ru'(t) + Sx'(t) + Ly'(t)) dt.$$

Proof. From (2.6), (2.9), (3.1) and (3.2) it follows (3.3). \square

We find conditions for the components of the vectors in T_pM , so that they are isotropic with respect to \tilde{g} .

Lemma 3.2. *Let $\{v, Qv, Q^2v, Q^3v\}$ be an orthonormal Q -basis of T_pM . If $r = xv + yQv + zQ^2v + uQ^3v$ is an isotropic vector, then its coordinates satisfy*

$$(3.4) \quad xz + yu = 0.$$

Proof. Taking into account (2.2), (2.6) and (2.7) we get (3.4). \square

Furthermore, due to Lemma 3.2, we obtain

Proposition 3.3. *If the vector force field (3.1) is isotropic, then its components satisfy*

$$(3.5) \quad PS + RL = 0.$$

Proposition 3.4. *If the curve (3.2) is isotropic, then the components of its tangent vector satisfy*

$$(3.6) \quad dx dz + dy du = 0.$$

Now, we suppose that F and c are both isotropic. Hence, having in mind (3.5) and (3.6), we consider the following cases.

Case (i) We assume that $dy = 0$ and $dx = 0$. Therefore, using (3.3), we obtain

$$(3.7) \quad A = \int_{\alpha}^{\beta} (P(c_1, c_2, z, u)z' + R(c_1, c_2, z, u)u') dt,$$

where c_1 and c_2 are specific constants.

Case (ii) If $dy = 0$ and $dz = 0$, then (3.3) implies

$$(3.8) \quad A = \int_{\alpha}^{\beta} (S(x, c_3, c_4, u)x' + R(x, c_3, c_4, u)u') dt,$$

where c_3 and c_4 are specific constants.

Case (iii) We suppose that $R = 0$ and $P = 0$. In this case equality (3.3) yields

$$(3.9) \quad A = \int_{\alpha}^{\beta} (Sx' + Ly') dt.$$

Case (iv) If $R = 0$ and $S = 0$, then (3.3) implies

$$(3.10) \quad A = \int_{\alpha}^{\beta} (Ly' + Pz') dt.$$

Case (v) Let $R \neq 0$, $dy \neq 0$ be valid. From (3.5) and (3.6) we find respectively

$$(3.11) \quad L = -\frac{PS}{R}, \quad du = -\frac{dxdz}{dy}.$$

By virtue of (3.3) and (3.11) we have

$$(3.12) \quad A = \int_{\alpha}^{\beta} \frac{(RSx'y' - PSy'^2 + PRy'z' - R^2x'z')}{Ry'^2} dt.$$

Cases (i) – (v) we summarize in the next statement.

Theorem 3.5. *Let F be an isotropic vector force field and let c be an isotropic smooth curve. Therefore the work A , determined by (3.3), takes one of the following forms:*

- (i) A is (3.7), if $dx = dy = 0$;
- (ii) A is (3.8), if $dy = dz = 0$;
- (iii) A is (3.9), if $P = R = 0$;
- (iv) A is (3.10), if $R = S = 0$;
- (v) A is (3.12), if $dy \neq 0$ and $R \neq 0$.

4 The work in a 3-dimensional subspace of T_pM

Let the unit vector v induce a Q -basis of T_pM . Then v induces four different Q -bases of three vectors, which are $\{v, Qv, Q^2v\}$, $\{Qv, Q^2v, Q^3v\}$, $\{v, Q^2v, Q^3v\}$ and $\{v, Qv, Q^3v\}$. According to (2.3) and (2.4) all pyramids constructed on these bases are equal. Thus we will consider only one of them, the 3-dimensional subspace ε of T_pM , spanned by vectors v , Qv and Q^2v .

Lemma 4.1. *Let ε be a subspace of T_pM with a normalized basis $\{v, Qv, Q^2v\}$. The system of vectors $\{e_1, e_2, e_3\}$, determined by the equalities*

$$(4.1) \quad \begin{aligned} e_1 &= \frac{v + Q^2v}{2 \cos \frac{\phi}{2}}, & e_3 &= \frac{v - Q^2v}{2 \sin \frac{\phi}{2}}, \\ e_2 &= \frac{(-\cos \varphi)v + (1 + \cos \phi)Qv - (\cos \varphi)Q^2v}{\sqrt{(1 + \cos \phi)(1 + \cos \phi - 2 \cos^2 \varphi)}}, \end{aligned}$$

is an orthonormal basis of ε .

Proof. Taking into account (2.1), (2.3), (2.4) and (4.1) we obtain

$$g(e_1, e_1) = g(e_2, e_2) = g(e_3, e_3) = 1, \quad g(e_1, e_2) = g(e_2, e_3) = g(e_1, e_3) = 0.$$

□

Lemma 4.2. *Let \tilde{g} be the associated metric with g on (M, g, Q) . Let ε be a subspace of $T_p M$ with a basis $\{v, Qv, Q^2v\}$. Then the system of vectors $\{e_1, e_2, e_3\}$, determined by the equalities (4.1), satisfy*

$$(4.2) \quad \begin{aligned} \tilde{g}(e_1, e_1) = -\tilde{g}(e_3, e_3) = 1, \quad \tilde{g}(e_2, e_2) = b, \\ \tilde{g}(e_1, e_2) = \tilde{g}(e_2, e_3) = \tilde{g}(e_1, e_3) = 0, \end{aligned}$$

where

$$(4.3) \quad b = \frac{\cos \phi + \cos^2 \phi - 2 \cos^2 \varphi}{1 + \cos \phi - 2 \cos^2 \varphi}.$$

Proof. Taking into account (2.1), (2.3), (2.4), (2.6) and (4.1) we find (4.2) and (4.3). \square

We consider a coordinate system p_{xyz} such that the vectors e_1, e_2 and e_3 lie on the axes p_x, p_y and p_z , respectively. Hereof p_{xyz} is an orthonormal coordinate system in ε . In the subspace ε of $T_p M$, a vector force field F and a smooth curve c are determined as follows:

$$(4.4) \quad F(x, y, z) = P(x, y, z)e_1 + R(x, y, z)e_2 + S(x, y, z)e_3,$$

$$(4.5) \quad c : r(t) = x(t)e_1 + y(t)e_2 + z(t)e_3, \quad t \in [\alpha, \beta] \subset \mathbb{R}.$$

Theorem 4.3. *Let F be an arbitrary vector force field and let c be an arbitrary smooth curve in ε , determined by (4.4) and (4.5). Then the work A done by a force F , with respect to \tilde{g} , along the curve c is*

$$(4.6) \quad A = \int_{\alpha}^{\beta} (Px'(t) + bRy'(t) - Sz'(t)) dt.$$

Proof. From (2.6) and (4.2) it follows

$$\begin{aligned} \tilde{g}(F, dr) &= \tilde{g}(Pe_1 + Re_2 + Se_3, dx e_1 + dy e_2 + dz e_3) \\ &= P\tilde{g}(e_1, e_1)dx + R\tilde{g}(e_2, e_2)dy + S\tilde{g}(e_3, e_3)dz \\ &= Pdx + bRdy - Sdz \end{aligned}$$

Then, using (2.9), we get (4.6). \square

In the following statements we find conditions for the components of the vectors in ε , such that they are isotropic.

Lemma 4.4. *Let $\{e_1, e_2, e_3\}$ be an orthonormal basis of ε , determined by (4.1). If $r = xe_1 + ye_2 + ze_3$ is an isotropic vector, then its coordinates satisfy*

$$(4.7) \quad by^2 = z^2 - x^2.$$

Proof. From (2.6), (2.7) and (4.2) the proof follows. \square

Remark 4.1. Bearing in mind (4.7) we will consider only the isotropic vectors determined by conditions: $b > 0$ and $|z| > |x|$, or $b = 0$ and $|z| = |x|$, or $b < 0$ and $|z| < |x|$.

Furthermore, due to Lemma 4.4 and Remark 4.1, we immediately have

Proposition 4.5. Let $\{e_1, e_2, e_3\}$ be an orthonormal basis of ε , determined by (4.1). The vector force field (4.4) is isotropic if its components satisfy

$$(4.8) \quad bR^2 = S^2 - P^2.$$

In particular, if $b > 0$ then $|P| < |S|$, if $b = 0$ then $|P| = |S|$, if $b < 0$ then $|P| > |S|$.

Proof. The equalities (4.4) and (4.7) imply (4.8). \square

Proposition 4.6. Let $\{e_1, e_2, e_3\}$ be an orthonormal basis of ε , determined by (4.1). The curve (4.5) is isotropic if

$$(4.9) \quad b(dy)^2 = (dz)^2 - (dx)^2.$$

In particular, if $b > 0$ then $|dz| > |dx|$, if $b = 0$ then $|dx| = |dz|$, if $b < 0$ then $|dz| < |dx|$.

Evidently, taking into account (4.8) and (4.9) we find

$$(4.10) \quad b^2 R^2 (dy)^2 = (S^2 - P^2)(dz^2 - dx^2).$$

Let us denote

$$(4.11) \quad \delta = \arccos \sqrt{\frac{1}{2}(\cos \phi + \cos^2 \phi)},$$

where $0 < \phi < \frac{\pi}{2}$.

Therefore, having in mind (2.5), (4.3) and (4.11), we state the following

Theorem 4.7. Let F be an isotropic vector force field and let c be an isotropic smooth curve in ε . If $\varphi = \delta$ or $\varphi = \pi - \delta$, then the work A , determined by (4.6), takes one of the following forms:

$$(i) \quad A = 2 \int_{\alpha}^{\beta} P x'(t) dt, \text{ if } S dz = -P dx;$$

$$(ii) \quad A = 0, \text{ if } S dz = P dx.$$

Proof. From (4.3) we get $b = 0$. Consequently (4.8) and (4.9) imply $|P| = |S|$ and $|dz| = |dx|$. Therefore, using (4.6) we find conditions (i) and (ii). \square

In a similar way, with the help of (2.5), (4.10) and (4.11), we obtain the next statements.

Theorem 4.8. Let F be an isotropic vector force field and let c be an isotropic smooth curve in ε . If $\varphi \in (\frac{\phi}{2}, \delta) \cup (\pi - \delta, \pi - \frac{\phi}{2})$, $|S| < |P|$, $R > 0$ and $|dz| < |dx|$, then the work A , determined by (4.6), takes the form

$$A = \int_{\alpha}^{\beta} (P x'(t) - \sqrt{(S^2 - P^2)(z'^2 - x'^2)} - S z') dt.$$

Theorem 4.9. *Let F be an isotropic vector force field and let c be an isotropic smooth curve in ϵ . If $\varphi \in (\delta, \pi - \delta)$, $|S| > |P|$, $R > 0$ and $|dz| > |dx|$, then the work A , determined by (4.6), takes the form*

$$A = \int_{\alpha}^{\beta} (Px'(t) + \sqrt{(S^2 - P^2)(z'^2 - x'^2)} - Sz')dt.$$

5 Work in a 2-plane spanned on $\{v, Q^2v\}$

Now, we consider an arbitrary 2-plane $\epsilon = \{v, Q^2v\}$ in T_pM . Because of (2.1) we have that the structure Q^2 is almost product. Every 2-plane of the type $\{v, Q^2v\}$ in an arbitrary tangent space of an almost product manifold admits an orthonormal basis of the type $\{l, Q^2l\}$ (see [14] and [17]).

We suppose that $\{l, Q^2l\}$ is an orthonormal basis of ϵ . On ϵ we construct a coordinate system p_{xy} such that l is on the axis p_x and Q^2l is on the axis p_y . Hence p_{xy} is an orthonormal coordinate system. An arbitrary vector force field F and an arbitrary smooth curve c in ϵ are given as follows:

$$(5.1) \quad F(x, y, z) = P(x, y)l + R(x, y)Q^2l,$$

$$(5.2) \quad c : r(t) = x(t)l + y(t)Q^2l, \quad t \in [\alpha, \beta] \subset \mathbb{R}.$$

Theorem 5.1. *Let F be an arbitrary vector force field and let c be an arbitrary smooth curve in ϵ , determined by (5.1) and (5.2). Then the work A done by a force F , with respect to \tilde{g} , along the curve c is*

$$(5.3) \quad A = \int_{\alpha}^{\beta} (P(x, y)y'(t) + R(x, y)x'(t))dt,$$

Proof. Equalities (2.3), (2.6), (2.9), (5.1) and (5.2) imply (5.3). \square

In the following statements we find conditions for the components of the vectors in ϵ , such that they are isotropic.

Lemma 5.2. *Let $\{l, Q^2l\}$ be an orthonormal basis of the 2-plane ϵ . If $r = xl + yQ^2l$ is an isotropic vector, then its coordinates satisfy*

$$xy = 0.$$

Proof. With the help of (2.3), (2.6) and (2.7) we get the proof. \square

Immediately, by virtue of Lemma 5.2, we obtain the following propositions.

Proposition 5.3. *The vector force field (5.1) is isotropic if its components satisfy*

$$(5.4) \quad PR = 0.$$

Proposition 5.4. *The curve (5.2) is isotropic if*

$$(5.5) \quad dx dy = 0.$$

Due to Theorem 5.1 and the above propositions we obtain the next statement.

Theorem 5.5. *Let F be an isotropic vector force field and let c be an isotropic smooth curve in the 2-plane ϵ . Hence the work A , determined by (5.3), takes one of the following forms:*

$$(i) \quad A = 0, \text{ if } P = 0 \text{ and } dx = 0,$$

$$(ii) \quad A = 0, \text{ if } R = 0 \text{ and } dy = 0,$$

$$(iii) \quad A = \int_{\alpha}^{\beta} (P(c_1, y)y'(t))dt, \text{ if } R = 0 \text{ and } dx = 0,$$

$$(iv) \quad A = \int_{\alpha}^{\beta} (R(x, c_2)x'(t))dt, \text{ if } P = 0 \text{ and } dy = 0,$$

where c_1, c_2 are specific constants.

Proof. The proof follows by virtue of (5.3), (5.4) and (5.5). \square

6 Work in a 2-plane spanned on $\{v, Qv\}$

We note that the investigations in this section are based mainly on the results in [6].

We suppose that the unit vector v induces a Q -basis of T_pM . We consider the 2-plane $\eta = \{v, Qv\}$ in T_pM .

Lemma 6.1. *Let η be a 2-plane of T_pM with a normalized basis $\{v, Qv\}$. The system of vectors $\{e_1, e_2\}$, determined by the equalities*

$$(6.1) \quad e_1 = \frac{1}{2 \cos \frac{\varphi}{2}}(v + Qv), \quad e_2 = -\frac{1}{2 \sin \frac{\varphi}{2}}(v - Qv)$$

is an orthonormal basis of η .

Proof. From (2.3), (2.4) and (6.1) we have immediately $g(e_1, e_1) = g(e_2, e_2) = 1$ and $g(e_1, e_2) = 0$. \square

On η we construct a coordinate system p_{xy} with basis vectors e_1 on the axis p_x and e_2 on the axis p_y , determined by (6.1). Hence p_{xy} is an orthonormal coordinate system.

With the help of (2.6) and (6.1) we state the following

Lemma 6.2. *Let \tilde{g} be the associated metric with g on (M, g, Q) . Let η be a 2-plane of T_pM with a basis $\{e_1, e_2\}$, determined by (6.1). Then the vectors e_1 and e_2 satisfy*

$$(6.2) \quad \tilde{g}(e_1, e_1) = k_1, \quad \tilde{g}(e_2, e_2) = k_2, \quad \tilde{g}(e_1, e_2) = 0,$$

where

$$(6.3) \quad k_1 = \frac{\cos \phi + \cos \varphi}{1 + \cos \varphi}, \quad k_2 = \frac{\cos \phi - \cos \varphi}{1 - \cos \varphi}.$$

In the 2-plane η , a vector force field F and a smooth curve c are determined as follows:

$$(6.4) \quad F(x, y) = P(x, y)e_1 + R(x, y)e_2,$$

$$(6.5) \quad c : r(t) = x(t)e_1 + y(t)e_2, \quad t \in [\alpha, \beta] \subset \mathbb{R}.$$

Theorem 6.3. *Let F be an arbitrary vector force field and let c be an arbitrary smooth curve in η , determined by (6.4) and (6.5). Then the work A done by a force F , with respect to \tilde{g} , along the curve c is*

$$(6.6) \quad A = \int_{\alpha}^{\beta} (k_1 P(x, y)x'(t) + k_2 R(x, y)y'(t)) dt.$$

Proof. From (2.6) and (6.2) it successively follows

$$\begin{aligned} \tilde{g}(F, dr) &= \tilde{g}(Pe_1 + Re_2, dx e_1 + dy e_2) \\ &= k_1 P \tilde{g}(e_1, e_1) dx + k_2 R \tilde{g}(e_2, e_2) dy + (P dy + R dx) \tilde{g}(e_1, e_2) \\ &= k_1 P dx + k_2 R dy, \end{aligned}$$

and using (2.9) we obtain (6.6). \square

Farther, we find conditions for the components of the vectors in η , so that they are isotropic with respect to \tilde{g} .

Lemma 6.4. *Let $\{e_1, e_2\}$ be an orthonormal basis of the 2-plane η , determined by (6.1). If $r = xe_1 + ye_2$ is an isotropic vector, then its coordinates satisfy*

$$k_1 x^2 + k_2 y^2 = 0, \quad k_1 k_2 \leq 0.$$

Proof. By equalities (2.5), (2.6), (2.7), (6.2) and (6.3) we get the proof. \square

Immediately, by virtue of Lemma 6.4, we obtain the following propositions.

Proposition 6.5. *The vector force field (6.4) is isotropic if its components satisfy*

$$(6.7) \quad k_1 P^2 + k_2 R^2 = 0, \quad k_1 k_2 \leq 0.$$

Proposition 6.6. *The curve (6.5) is isotropic if*

$$(6.8) \quad k_1 (dx)^2 + k_2 (dy)^2 = 0, \quad k_1 k_2 \leq 0.$$

Let F be an isotropic vector force field and let c be an isotropic smooth curve in η . According to the values of the coefficients k_1 and k_2 we have two cases.

First we consider the case when $k_1 k_2 = 0$. If $\varphi = \phi \neq \frac{\pi}{2}$, then (6.3) implies $k_1 \neq 0$, $k_2 = 0$. Consequently, from (6.7) and (6.8) we have $P = 0$, $dx = 0$, which implies $A = 0$. Similarly, if we suppose that $\varphi = \pi - \phi$, then $A = 0$.

Therefore we obtain the next statement.

Theorem 6.7. *Let F be an isotropic vector force field and let c be an isotropic smooth curve in the 2-plane ϵ . If $\varphi = \phi$ or $\varphi = \pi - \phi$, then the work determined by (6.6) is $A = 0$.*

Now, we consider the case when $k_1 k_2 < 0$. Bearing in mind inequalities (2.5), we have that both angles ϕ and φ satisfy one of the following conditions:

$$(6.9) \quad \phi \in (0, \frac{\pi}{2}), \quad \varphi \in (\frac{\phi}{2}, \phi) \cup (\pi - \phi, \pi - \frac{\phi}{2});$$

$$(6.10) \quad \phi \in (\frac{\pi}{2}, \frac{2\pi}{3}), \quad \varphi \in (\frac{\phi}{2}, \pi - \phi) \cup (\phi, \pi - \frac{\phi}{2});$$

$$(6.11) \quad \phi = \frac{\pi}{2}, \quad \varphi \neq \frac{\phi}{2}.$$

In this case, the curve $c : k_1 x^2 + k_2 y^2 = 0$ degenerates into two lines with equations:

$$(6.12) \quad y = \pm \sqrt{\frac{-k_1}{k_2}} x, \quad k_1 k_2 < 0.$$

The above considerations imply the next statement.

Theorem 6.8. *Let F be an isotropic vector force field and let c be an isotropic smooth curve in the 2-plane ϵ . If one of the conditions (6.9) – (6.11) is valid, then the work determined by (6.6) takes one of the following forms:*

$$(i) \quad A = 0, \text{ if } R = \pm \sqrt{\frac{-k_1}{k_2}} P, \text{ } dy = \mp \sqrt{\frac{-k_1}{k_2}} dx;$$

$$(ii) \quad A = 2 \int_{\alpha}^{\beta} P(t, \pm \sqrt{\frac{-k_1}{k_2}} t) dt, \text{ if } R = \pm \sqrt{\frac{-k_1}{k_2}} P \text{ and } dy = \pm \sqrt{\frac{-k_1}{k_2}} dx.$$

Proof. Substituting (6.12) into (6.6) we get conditions (i) and (ii). \square

References

- [1] C. L. Bejan and S. L. Druță-Romanuic, *Walker manifolds and Killing magnetic curves*, Differential Geom. Appl. 35 (2014), 106–116.
- [2] J. Del Amor, A. Giménez and P. Lucas, *Null curve evolution in four-dimensional pseudo-Euclidean spaces*, Adv. Math. Phys. (2016), Art. ID 5725234, 15 pp.
- [3] I. Dokuzova, *Curvature properties of 4-dimensional Riemannian manifolds with a circulant structure*, J. Geom. 108, 2 (2017), 517–527.
- [4] I. Dokuzova, *Almost conformal transformation in a four-dimensional Riemannian manifold with an additional structure*, Int. Electron. J. Geom. 6, 2 (2013), 9–13.
- [5] I. Dokuzova, *Four-dimensional Riemannian product manifolds with circulant structures*, (2019), arXiv:1904.09424.
- [6] I. Dokuzova and G. Dzhelapov, *Spheres and circles in a tangent space of a 4-dimensional Riemannian manifold with respect to an indefinite metric*, J. Geom. (2018), 109:49.
- [7] I. Dokuzova, D. Razpopov and M. Manev, *Two types of Lie groups as 4-dimensional Riemannian manifolds with circulant structure*, In: Maths. Educ. Math. Proc. 47th Spring Conf. UBM, Borovets, (2018), 115–120.
- [8] K. L. Duggal and A. Bejancu, *Lightlike submanifolds of semi-Riemannian manifolds and applications*. Mathematics and its Applications 364, Kluwer Academic Publishers Group, Dordrecht, 1996. 300 pp.

- [9] K. L. Duggal and D. H. Jin, *Null curves and hypersurfaces of semi-Riemannian manifolds*. World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2007. 293 pp.
- [10] D. Gribacheva and D. Razpopov, *Riemannian almost product manifolds generated by a circulant structure*, J. Tech. Univ. Plovdiv, Fundam. Sci. Appl., Ser. A, Pure Appl. Math. 23 (2017), 171–174.
- [11] A. Kazan and H.B. Karadağ, *Magnetic pseudo null and magnetic null curves in Minkowski 3-space*, International Mathematical Forum 12, 3 (2017), 119-132.
- [12] T. Körpınar and R. C. Demirkol, *Energy on a timelike particle in dynamical and electrodynamic force fields in De-Sitter space*, Rev. Mexicana Fis. 63, 6 (2017), 560-568.
- [13] H. Manev and G. Nakova *Slant null curves on normal almost contact B-metric 3-manifolds with parallel reeb vector field*, Results Math., **71** (2017), 933-947.
- [14] D. Mekerov, *On Riemannian almost product manifolds with nonintegrable structure*, J. Geom. 89, 1-2 (2008), 119–129.
- [15] Z. Özdemir, *Magnetic trajectories on lightlike surfaces*, Differ. Equ. Dyn. Syst. (2020), <https://doi.org/10.1007/s12591-020-00519-7>.
- [16] D. Razpopov, *Four-dimensional Riemannian manifolds with two circulant structures*, In: Maths. Educ. Math. Proc. of 44-th Spring Conf. of UBM, SOK Kamchia, Bulgaria (2015), 179–185.
- [17] M. Staikova, K. Gribachev and D. Mekerov, *Invariant hypersurfaces of Riemannian P-manifolds*, Plovdiv. Univ. Paisii Khilendarski Nauchn. Trud. Mat. 25, 3 (1987), 253–266 (in Bulgarian)

Authors' address:

Dimitar Razpopov, Georgi Dzhelepov
Department of Mathematics and Informatics,
Agricultural University of Plovdiv,
12 Mendeleev Blvd., 4000 Plovdiv, Bulgaria.
E-mail addresses: razpopov@au-plovdiv.bg , dzhelepov@abv.bg